

METHODS, PRODUCTS AND TREATMENTS FOR DIABETES

RELATED APPLICATION

This application claims priority under 35 U.S.C. §119 from U.S. provisional
5 application serial number 60/203,254 filed May, 8, 2000.

BACKGROUND OF THE INVENTION

Diabetes Mellitus (diabetes) is a leading cause of morbidity and mortality in the
adult population. This is primarily because diabetic patients tend to develop vascular
10 complications that involve the kidneys (diabetic nephropathy), the retina (diabetic
retinopathy), as well as large and small blood vessels in other organs (macro- and micro-
vascular disease) including nerves (diabetic neuropathy). It is well established that the
vascular complications of diabetes are caused by elevated blood glucose levels over long
periods of time. Elevated blood glucose levels affect proteins by a process known as
15 glycation. Different "glycated" proteins have been identified in diabetic subjects,
including albumin, hemoglobin and others. Measurement of the extent of protein
"glycation" of certain proteins is considered a valuable clinical tool to assess long term
glycemic control and thereby the efficacy of diabetes treatment.

Glycation, the non-enzymatic attachment of glucose to proteins, is considered a
20 major pathophysiological mechanism causing tissue damage in diabetic subjects.
Glycation involves the reaction of glucose and/or other reducing sugars with amino
groups in proteins resulting in the formation of a Schiff base or aldimine. This labile
adduct can tautomerize via the Amadori rearrangement to the more stable ketoamine.
The function of the glycated protein may be impaired, depending on the location of the
25 amino group(s) affected. For example, amino-terminal glycation of the β -chains of
hemoglobin gives rise to the glycated hemoglobins (HbA1) in which responsiveness to
2,3-diphosphoglycerate is decreased and oxygen affinity increased. Glycation of the
major thrombin inhibitor of the coagulation system, antithrombin III, decreases its
affinity for heparin, and has been postulated to contribute to the hypercoagulable state
30 associated with diabetes.

Hemoglobin glycation and thrombin inhibitor glycation do not account for the vascular complications of diabetes. The mechanism which results in such complications remains unknown.

Currently, protein glycation in diabetic subjects is measured in blood by
5 estimating the amount of glycated hemoglobin (hemoglobin A1c) through a complicated clinical test that requires extraction of a blood sample. Accordingly, there is a need for a simplified and less invasive method for rapid monitoring of protein glycation levels.

SUMMARY OF THE INVENTION

The glycation of the membrane protein known as CD59, a key regulator of the
10 complement system, has been discovered, according to the invention, as being involved in the pathogenesis of the vascular complications of diabetes. Unexpectedly, glycated CD59 has now been identified in human urine. The amount of glycated CD59 in human urine correlates with glycemic control and levels of glycated hemoglobin. Detection of glycated CD59 in the urine of diabetic patients can be used to monitor glycemic control
15 in diabetic patients, and to select subjects for therapy.

The invention, in one aspect, is a method of characterizing the impact of blood sugar levels on abnormal glycation levels in a subject. The method involves obtaining a level of the amount of K41-glycated CD59 from a sample obtained from the subject.

In one embodiment the method involves comparing the level of K41-glycated
20 CD59 to a control. In one embodiment, the sample is a fluid sample. Preferred samples are urine and blood, most preferably, urine. In one embodiment the subject is diabetic.

The subject can be receiving treatment for regulating blood sugar levels.

In one embodiment the level of K41-glycated CD59 is obtained using an agent that binds CD59. Preferably, the agent binds specifically CD59. Most preferably, the
25 agent binds specifically K41-glycated CD59. The level of K41-glycated CD59 can be obtained, for example, using an immunoassay.

The invention in another aspect is a method of evaluating a treatment for regulating blood sugar levels. The method includes obtaining a first level of the amount of K41-glycated CD59 from a first sample obtained from a subject undergoing the
30 treatment for regulating blood sugar levels, and obtaining a second level of the amount of K41-glycated CD59 from a second sample obtained from the subject after obtaining the

first level, and comparing the first level to the second level as an indication of evaluation of the treatment. Preferably, the second level is obtained at least one day after the first level. In important embodiments, the second level is a plurality of second levels spaced out over time so that long term treatment can be evaluated and monitored.

5 In one embodiment the subject is diabetic. The subject can be undergoing treatment with an oral blood sugar regulating agent. In another embodiment, the subject is undergoing treatment with insulin or an insulin analog.

 The samples preferably are fluid samples. The samples can be urine or blood, preferably urine. In one embodiment the levels are obtained using an immunoassay.

10 The invention in another aspect involves a method of selecting a treatment for regulating blood sugar levels in a subject. The method includes obtaining a level of the amount of K41-glycated CD59 from a sample obtained from the subject, and selecting the treatment for regulating blood sugar levels in the subject based at least in part on the level obtained. In one embodiment the subject is diabetic. The subject can already be
15 receiving drug therapy for regulating blood sugar levels. In another embodiment, the subject has not yet received drug therapy for regulating blood sugar levels.

 Preferred embodiments, including samples and assays, are as described above.

 Another aspect of the invention is a method for determining regression, progression or onset of a condition characterized by abnormal levels of glycated protein.

20 The method involves obtaining a level of the amount of K41-glycated CD59 from a sample obtained from a subject, and comparing the level to a control as a determination of regression, progression or onset of the condition. The sample can be a fluid sample. Preferably, the fluid sample is blood or urine, most preferably urine.

 The subject can be diabetic. In one embodiment, the subject is undergoing drug
25 therapy for regulating blood sugar levels.

 The level can be obtained using an agent that binds to K41-glycated CD59. Preferably, the agent binds specifically to K41-glycated CD59. In one embodiment the level is obtained using an immunoassay.

 Another aspect of the invention is a method of treating a subject to reduce the risk
30 of or the progression of a disorder associated with abnormally high levels of K41-glycated CD59. The method of treatment involves selecting and administering to a subject who is known to have an abnormally high level of K41-glycated CD59 an agent

for reducing K41-glycated CD59 levels in an amount effective to reduce K41-glycated CD59 levels. In one embodiment the subject is diabetic. In one embodiment the agent is an oral blood sugar regulating agent and the subject is otherwise free of symptoms calling for any oral blood sugar regulating agent. In one embodiment the agent is insulin
5 or an insulin analog and the subject is otherwise free of symptoms calling for treatment with insulin.

In one embodiment the subject already is being treated with a therapeutic for lowering blood sugar levels and the therapeutic is not the agent. In another embodiment the subject already is being treated with the agent in a previous amount and wherein the
10 effective amount is a higher dose than the previous amount.

Another aspect of the invention is a composition of matter that is isolated K41-glycated CD59. In one embodiment the isolated K41-glycated CD59 represents at least 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or 95% of CD59 in the composition.

Another aspect of the invention is a composition of matter that is pure K41-glycated CD59 or a fragment thereof, wherein the fragment comprises at least 8, 9, 10,
15 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 contiguous amino acids having a consecutive sequence found in CD59 and including K41, and wherein K41 is glycated.

Another aspect of the invention is an agent that binds specifically to K41-glycated CD59 but not to K41-nonglycated CD59. In one embodiment the agent is an
20 antibody.

According to still another aspect of the invention, a kit is provided for determining the level of K41-glycated CD59 and a sample. The kit comprises a package containing (1) an agent that binds specifically to only one of K41-glycated CD59 and K41-nonglycated CD59, and (2) a control. In one embodiment, the agent is two agents, a
25 first agent that binds both glycated and nonglycated CD59 and a second agent that binds only one of a glycated K41 and a nonglycated K41. Thus, the first agent can bind specifically CD59, without distinguishing K41-glycated and K41-nonglycated CD59, and the second agent may be capable of distinguishing a glycated and a nonglycated lysine motif. In a preferred embodiment, the agent binds specifically K41-glycated
30 CD59 or K41 nonglycated CD59. In one preferred embodiment, the agent is an antibody or a fragment of an antibody. The kit may further comprise instructions for using the kit for any of the methods described above. In another embodiment, the agent is a first

agent and the kit further comprises a second agent. The first agent binds specifically to K41-glycated CD59 and the second agent binds specifically to K41-nonglycated CD59.

These and other aspects of the invention will be described in greater detail below.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG 1 is a schematic of a kit according to the invention.

FIG. 2 is a graph indicating CD59 activity as a percent of control activity for a cell suspension of GPE incubated with purified CD59 previously exposed for different time intervals to the glyating sugars glucose (open triangles, 0.5M) or ribose (open squares; 0.5M), or to non-glycating sorbitol (closed squares, 0.5M). Data points represent the mean of triplicate determinations (SEM are smaller than data points). Figure is representative of experiments run three times with comparable results. Inset shows the number of CD59 molecules incorporated per GPE as determined with ^{125}I -CD59 before (C) and after (R) glycation with ribose.

FIG. 3A is an image of a western blot and FIGs 3B and 3C are graphs. Panel (3A) indicates expression of wild-type (WT) CD59, and CD59 mutants Gln-41 and Gln-44 in CHO cells as confirmed by SDS-PAGE followed by Western blot analysis of CHO cell extracts using the YTH 53.1 monoclonal antibody, Vector (closed square); WT (open square); CD59-Gln-41 (open triangle); Gln-44 (open circle). Panel (3B) indicates activity of WT and mutant CD59s tested in the GPE hemolytic assay before and after glycation with ribose for the time intervals indicated. Panel (3C) indicates CD59 and mutant CD59 activity as a percent of control activity for a cell suspension of GPE incubated with purified WT CD59 (open squares) and Gln-41 (open triangles), and Gln-44 (open circles) following incubation with the glyating sugar ribose. Data points represent the mean of triplicate determinations (SEM are smaller than the data points). The figure is representative of experiments run three times with comparable results.

FIG 4 A and 4B are bar graphs. Panel (4A) Bar graph indicating HUVEC incubated with or without 50 mM ribose followed by reduction with cyanoborohydride. Cells were then exposed to purified terminal complement components C5b6, C7, C8 and C9 to form the MAC and the mitogenic activity in the conditioned medium was measured. Results are expressed as the ratio of mitogenic activity released into the conditioned media in the presence or absence of MAC. Panel (4B). Indicates Human

RBC incubated without or with 50 mM ribose followed by reduction with cyanoborohydride to stabilize labile Schiff base adducts. After volume adjustment, aliquots of glycated and non glycated cells were separated for rescue with purified CD59 (4μg), exposure to neutralizing anti-CD59 antibody (YTH53.1) or both, and then
5 exposed to purified C5b6, C7, C8 and C9 to form the MAC. Open columns represent control cells not exposed to cyanoborohydride, filled columns represent cells exposed to cyanoborohydride, as above.

FIG 5A and 5B are images of immunoblots. Panel (5A) is an immunoblot of urine samples from non-diabetic (lanes 1,4 and 5) and diabetic (lanes 2 and 3) subjects.
10 Urine was concentrated by ultrafiltration, separated by anion exchange chromatography and fractions dot-blotted for the presence of CD59 with anti-CD59 specific antibody. CD59-positive fractions were pooled and immunoprecipitated with the HC1 anti-CD59 specific antibody. The immunoprecipitate was separated by SDS-PAGE and immunoblotted. Panel (5A), upper blot: immunoblot of CD59 positive fractions of the immunoprecipitate with the monoclonal anti-CD59 YTH53.1. Lower blot: The immunoprecipitate separated by SDS-PAGE and immunoblotted with the anti-hexitol-lysine antibody . Lower section, Panel A indicates levels of glycated CD59 and glycated hemoglobin (HbA1c). Panel (5B). Immunoblot of glycated (G) and non-glycated (NG) albumin (left) and affinity chromatography purified CD59 (right) with anti-hexitol-lysine
15 antibody.
20 antibody.

DETAILED DESCRIPTION

The invention disclosed herein describes novel methods and compositions for detecting and measuring K41-glycated CD59 levels, particularly as they relate to glycemic levels. The discovery that CD59 is glycated facilitates analysis of diseases in
25 which the amount of CD59 glycation differs from normal levels. For example, it has been discovered that the level of glycation of CD59 is elevated in diabetes. Thus, onset, progression and/or regression of diabetes or other diseases can be monitored by monitoring levels of glycated CD59 in a subject. It also has been determined that CD59 is present in urine. Therefore, the measurement can be done in urine without requiring a
30 blood sample.

In addition, in contrast to markers of glycation such as hemoglobin, glycation of CD59 is believed to be involved in the pathogenesis of the vascular complications of

diabetes. Accordingly, clinical evaluation of K41-glycated CD59 in urine is a direct measure for vascular damage induced by glycation. Without wishing to be bound by any particular theory, it is believed that patients with abnormally high levels of K41-glycated CD59 in urine will either have or will be more prone to develop vascular complications of diabetes.

As used herein, CD59 (also known as membrane inhibitor of reactive lysis [MIRL], protectin, HRF20 and H19) and K41-glycated CD59 are a polypeptide having substantial nucleotide and/or amino acid sequence identity of Accession No. M95708 (Davies, A., et al., *Journal J. Exp. Med.* 170 (3), 637-654 (1989)).

As used herein, "K41-glycated CD59" means CD59 which has been glycated at amino acid No. 41 of CD59, which is a lysine (hence K41). It has been determined, surprisingly, that K41-glycation of CD59 is correlated to abnormal blood sugar levels and that glycation at K41 interferes with the normal activity of CD59. CD59 functions normally by binding to the terminal components of the membrane attack complex of complement (MAC), thereby interfering with membrane insertion and polymerization of the C9 component of complement. Glycation at the K41 of CD59 interferes with CD59's ability to prevent the assembly of the MAC. It is believed that, as a result of K41-glycation of CD59, the MAC is permitted to be activated and leads to the development of proliferative chronic diabetic complications. Indeed, the present inventor has shown that the membrane attack complex stimulates proliferation of fibroblasts, smooth muscle, mesangial and other cells, in part by releasing growth factors such as FGF and PDGF from MAC-targeted endothelium. The MAC also induces increased synthesis of extracellular matrix proteins by mesangial cells. Thus, increased MAC deposition in diabetic tissues is believed to induce growth factor release from endothelium, which stimulates cell proliferation in the vascular wall and contributes to the expansion of the extracellular matrix and to the glomerulosclerosis that characterizes diabetic nephropathy.

Thus, according to one aspect of the invention, isolated K41-glycated CD59 is provided. By "isolated", it is meant separated from its native environment and present in sufficient quantity to permit its identification or use according to the procedures described herein. Isolated includes (1) selectively produced by expression cloning and glycation or (2) purified as by immunoprecipitation, chromatography or electrophoresis.

Because an isolated material may be admixed with a carrier in a preparation, such as, for example, for injecting into an animal, the isolated material may comprise only a small percentage by weight of the preparation. The material is none-the-less isolated in that it has been separated from the substances with which it typically is associated in living systems, i.e. from K41-nonglycated CD59. According to the present invention, the isolated K41-glycated CD59 represents at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or 95% of the CD59 in the composition (i.e. percentage of total CD59 both K41-glycated and K41-nonglycated).

According to another aspect of the invention, pure K41-glycated CD59 is provided. Isolated proteins or polypeptides may, but need not be, pure. The term "pure" means the proteins or polypeptides are essentially free of other substances with which they may be found in nature or in *in vivo* systems to an extent practical and appropriate for their intended use. Pure polypeptides may be produced by techniques well-known in the art. As used herein, "pure" means at least 95% of the total CD59 is K41-glycated CD59.

The invention also involves fragments of the foregoing proteins. A fragment of K41-glycated CD59 comprises at least 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 contiguous amino acids of CD59 having a consecutive sequence found in CD59 and including K41, wherein K41 is glycated. Fragments of K41-glycated CD59 can be used for a variety of purposes, including in the preparation of molecules that bind specifically to K41-glycated CD59 and in immunoassays well-known to those of ordinary skill in the art, including competitive binding immunoassays. One such peptide comprises amino acids 36-50 of CD59, wherein the N-terminus and K38 are both blocked to prevent glycation at these points. Virtually any fragment, including the active site of CD59 is useful according to the invention. Fragments including (N-term) YNKAWKFEHANFNDC (C term) SEQ ID NO: 1 are particularly useful according to the invention for raising antibodies to glycated CD59. Thus, as will be apparent to those of ordinary skill in the art, fragments beginning anywhere from amino acid No. 1, 2, 3...(to and including amino acid No. 36) and peptides ending at amino acid 50, 51, 52, 53...(to and including amino acid 77) are particularly useful according to the invention, for human sequences. As is well-known to those of ordinary skill in the art, sequences of non-human subjects differ.

As mentioned above, by having isolated or pure material, or fragments thereof, agents can be prepared that bind specifically to K41-glycated CD59. As used herein, “binding specifically to” means capable of distinguishing the identified material from other materials sufficient for the purpose to which the invention relates. Thus, “binding specifically to” CD59 means the ability to bind to and distinguish CD59 from other proteins. Binding specifically to K41-glycated CD59 means binding to and distinguishing K41-glycated CD59 from K41-nonglycated CD59.

Agents that bind to CD59 molecules and/or K41-glycated CD59 molecules, and fragments thereof, include polypeptide agents. Such polypeptides include polyclonal and monoclonal antibodies, prepared according to conventional methodology. Antibodies that bind CD59 molecules and/or K41-glycated CD59 molecules and are useful for determining glycated CD59 levels, include but are not limited to: antibodies that bind specifically to CD59, antibodies that bind specifically to fragments of CD59, antibodies that bind specifically to K-glycated regions of proteins including CD59, and antibodies that bind specifically to K41-glycated CD59 or fragments of CD59. Certain antibodies useful in the methods of the invention already are known in the art and include anti-CD59 antibodies as well as anti-glycation-moiety antibodies, for example Anti-CD-59 YTH53.1, and the anti-hexitol-lysine antibody, which binds to glycated CD59. Agents that bind specifically K41-glycated CD59 but not K41-nonglycated CD59, and vice-versa, are believed novel and part of the present invention.

Significantly, as is well-known in the art, only a small portion of an antibody molecule, the paratope, is involved in the binding of the antibody to its epitope (see, in general, Clark, W. R. (1986) *The Experimental Foundations of Modern Immunology*, Wiley & Sons, Inc., New York; Roitt, I. (1991) *Essential Immunology*, 7th Ed., Blackwell Scientific Publications, Oxford). The pFc' and Fc regions, for example, are effectors of the complement cascade but are not involved in antigen binding. An antibody from which the pFc' region has been enzymatically cleaved, or which has been produced without the pFc' region, designated an F9ab').sub.2 fragment, retains both of the antigen binding sites of an intact antibody. Similarly, an antibody from which the Fc region has been enzymatically cleaved, or which has been produced without the Fc region, designated an Fab fragment, retains one of the antigen binding sites of an intact antibody molecule. Proceeding further, Fab fragments consist of a covalently bound

antibody light chain and a portion of the antibody heavy chain denoted Fd. The Fd fragments are the major determinant of antibody specificity (a single Fd Fragment may be associated with up to ten different light chains without altering antibody specificity) and Fd fragments retain epitope-binding ability in isolation.

5 Within the antigen-binding portion of an antibody, as is well-known in the art, there are complementarity determining regions (CDRs), which directly interact with the epitope of the antigen, and framework regions (Frs), which maintain the tertiary structure of the paratope (see, in general, Clark, W. R. (1986) *The Experimental Foundations of Modern Immunology*, Wiley & Sons, Inc., New York; Roitt, I. (1991) *Essential*
10 *Immunology*, 7th Ed., Blackwell Scientific Publications, Oxford). In both the heavy chain Fd fragment and the light chain of IgG immunoglobulins, there are four framework regions (FR1 through FR4) separated respectively by three complementarity determining regions (CDR1 through CDR3). The CDRs, and in particular the CDR3 regions, and more particularly the heavy chain CDR3, are largely responsible for antibody specificity.

15 It is now well-established in the art that the non-CDR regions of a mammalian antibody may be replaced with similar regions of conspecific or heterospecific antibodies while retaining the epitopic specificity of the original antibody. This is most clearly manifested in the development and use of "humanized" antibodies in which non-human CDRs are covalently joined to human FR and/or Fc/pFc' regions to produce a functional
20 antibody. See, e.g., U.S. patents 4,816,567, 5,225,539, 5,585,089, 5,693,762 and 5,859,205.

 Thus, for example, PCT International Publication Number WO 92/04381 teaches the production and use of humanized murine RSV antibodies in which at least a portion of the murine FR regions have been replaced by FR regions of human origin. Such
25 antibodies, including fragments of intact antibodies with antigen-binding ability, are often referred to as "chimeric" antibodies.

 Thus, as will be apparent to one of ordinary skill in the art, the present invention also provides for F(ab').sub.2, Fab, Fv and Fd fragments; chimeric antibodies in which the Fc and/or Fr and/or CDR1 and/or CDR2 and/or light chain CDR3 regions have been
30 replaced by homologous human or non-human sequences; chimeric F(ab').sub.2 fragment antibodies in which the FR and/or CDR1 and/or CDR2 and/or light chain CDR3 regions have been replaced by homologous human or non-human sequences;

chimeric Fab fragment antibodies in which the FR and/or CDR1 and/or CDR2 and/or light chain CDR3 regions have been replaced by homologous human or non-human sequences; and chimeric Fd fragment antibodies in which the FR and/or CDR1 and/or CDR2 regions have been replaced by homologous human or nonhuman sequences. The present invention also includes so-called single chain antibodies.

Thus, the invention involves polypeptides of numerous size and type that bind specifically to K41-glycated or K41-nonglycated CD59 molecules. These polypeptides may be derived also from sources other than antibody technology. For example, such polypeptide binding agents can be provided by degenerate peptide libraries which can be readily prepared in solution, in immobilized form or as phage display libraries. Combinatorial libraries also can be synthesized of peptides containing one or more amino acids. Libraries further can be synthesized of peptoids and non-peptide synthetic moieties.

Phage display can be particularly effective in identifying binding peptides useful according to the invention. Briefly, one prepares a phage library (using e.g. m13, fd, or lambda phage), displaying inserts from 4 to about 80 amino acid residues using conventional procedures. The inserts may represent, for example, a completely degenerate or biased array. One then can select phage-bearing inserts which bind to K41-glycated or K41-nonglycated CD59 molecules. This process can be repeated through several cycles of reselection of phage that bind to the CD59 molecules. Repeated rounds lead to enrichment of phage bearing particular sequences. DNA sequences analysis can be conducted to identify the sequences of the expressed polypeptides. The minimal linear portion of the sequence that binds to the CD59 molecules can be determined. One can repeat the procedure using a biased library containing inserts containing part of all of the minimal linear portion plus one or more additional degenerate residues upstream or downstream thereof. Yeast two-hybrid screening methods also may be used to identify polypeptides that bind to the CD59 molecules. Thus, K41-glycated CD59 molecules and K41-nonglycated CD59 molecules, or fragments thereof containing the glycation site, can be used to screen peptide libraries, including phage display libraries, to identify and select peptide binding partners of the K41-glycated or K41-nonglycated CD59 molecules.

As detailed herein, the foregoing antibodies and other binding molecules may be used for example to isolate and identify CD59 protein and/or K41-glycated and/or K41-nonglycated CD59 protein. The antibodies may be coupled to specific diagnostic labeling agents for imaging of the protein or fragment thereof. The antibodies may also be used for immunoprecipitation, immunoblotting CD59 and/or K41-glycated CD59 using standard methods known to those of ordinary skill in the art.

The invention also involves a variety of assays based upon detecting levels of K41-glycated CD59 in subjects. The assays include (1) characterizing the impact of blood sugar levels on glycation levels in a subject; (2) evaluating a treatment for regulating blood sugar levels in a subject; (3) selecting a treatment for regulating blood sugar levels in a subject; and (4) determining progression, progression or onset of a condition characterized by abnormal levels of glycated protein in a subject. Thus, subjects can be characterized, treatment regimens can be monitored, treatments can be selected and diseases can be better understood using the assays of the present invention. For example, the invention provides in one aspect a method for measuring the level of K41-glycated CD59 in a subject, which is a direct indicator of the level of the subject's glycemic control. The impact of blood sugar levels or glycation levels thus can be measured due to the positive correlation between the level of circulating blood glucose and the amount of glycation of endogenous CD59. The level of K41-glycated CD59 thus correlates with the level of glycemic control in the subject. Relatively low levels of K41-glycated CD59 reflect well-controlled circulating blood sugar levels and selectively high levels of K41-glycated CD59 reflect poorly controlled glycemic levels.

The assays described herein are carried out on samples obtained from subjects. As used herein, a subject is a human, non-human primate, cow, horse, pig, sheep, goat, dog, cat, or rodent. In all embodiments, human subjects are preferred. The samples used herein are any body tissue or body fluid sample obtained from a subject. Preferred are body fluids, for example, lymph, saliva, blood, urine, and the like. Blood and urine are preferred, urine being most preferred. It has been surprisingly discovered that glycated CD59 can be detected in urine, thereby obviating the need for a blood sample.

Particularly, important subjects to which the present invention can be applied are diabetic subjects.

The term "diabetic" as used herein, means an individual who, at the time the sample is taken, has a primary deficiency of insulin. The term diabetic includes, but is not limited to, individuals with juvenile diabetes (Type 1 diabetes), adult-onset diabetes (Type 2 diabetes), gestational diabetes, and any other conditions of insulin deficiency.

5 The term "diabetic" is a term of art, known and understood by those practicing in the medical profession, a formal definition of which can be found in Harrison's Principles of Medicine (Harrisons, Vol 14, Principles of Internal Medicine, Eds. Fauci, A.S., E. Braunwald, K.J. Isselbacher, J.D. Wilson, J.B. Martin, D.L. Kasper, S.L. Hauser, D.L. Longo, McGraw-Hill, New York, 1999).

10 All of the assays described herein involve measuring levels of K41-glycated CD59. Levels of K41-glycated CD59 can be determined in a number of ways when carrying out the various methods of the invention. In one particularly important measurement, the level of K41-glycated CD59 is measured in relation to K41-nonglycated CD59. Thus, the measurement is a relative measure, which can be
15 expressed, for example, as a percentage of total CD59. Another measurement of the level of K41-glycated CD59 is a measurement of absolute levels of K41-glycated CD59. This could be expressed, for example, in terms of grams per liter of body fluid. Another measurement of the level of K41-glycated CD59 is a measurement of the change in the level of K41-glycated CD59 over time. This may be expressed in an absolute amount or
20 may be expressed in terms of a percentage increase or decrease over time.

Importantly, levels of K41-glycated CD59 are advantageously compared to controls according to the invention. The control may be a predetermined value, which can take a variety of forms. It can be a single cut-off value, such as a median or mean. It can be established based upon comparative groups, such as in groups having normal
25 amounts of circulating insulin and groups having abnormal amounts of circulating insulin. Another example of comparative groups would be groups having a particular disease, condition or symptoms and groups without the disease, condition or symptoms. Another comparative group would be a group with a family history of a condition and a group without such a family history. The predetermined value can be arranged, for
30 example, where a tested population is divided equally (or unequally) into groups, such as a low-risk group, a medium-risk group and a high-risk group or into quadrants or quintiles, the lowest quadrant or quintile being individuals with the lowest risk or

amounts of glycated protein and the highest quadrant or quintile being individuals with the highest risk or amounts of glycated protein.

The predetermined value, of a course, will depend upon the particular population selected. For example, an apparently healthy population will have a different 'normal' range than will a population which is known to have a condition related to abnormal protein glycation. Accordingly, the predetermined value selected may take into account the category in which an individual falls. Appropriate ranges and categories can be selected with no more than routine experimentation by those of ordinary skill in the art. By abnormally high it is meant high relative to a selected control. Typically the control will be based on apparently healthy normal individuals in an appropriate age bracket.

In measuring the relative amount of K41-glycated CD59 to K41-nonglycated CD59, those of ordinary skill in the art will appreciate that the relative amount may be determined by measuring either the relative amount of K41-glycated CD59 or the relative amount of K41-nonglycated CD59. In other words, if 90% of an individual's CD59 is K41-nonglycated CD59, then 10% of the individual's CD59 will be K41-glycated CD59. Thus, measuring the level of K41-glycated CD59 may be carried out by measuring the relative amount of K41-nonglycated CD59.

It will also be understood that the controls according to the invention may be, in addition to predetermined values, samples of materials tested in parallel with the experimental materials. Examples include samples from control populations or control samples generated through manufacture to be tested in parallel with the experimental samples.

The various assays used to determine the levels of K41-glycated CD59 include: binding assays, such as described in the examples below; specific binding assays, such as using materials which bind specifically to K41-glycated CD59; gel electrophoresis; NMR; and the like. Immunoassays may be used according to the invention including sandwich-type assays, competitive binding assays, one-step direct tests and two-step tests such as described in the examples.

As mentioned above, it is also possible to characterize blood sugar levels by monitoring changes in the absolute or relative amounts of glycated CD59 over time. For example, it is expected that an increase in glycated CD59 correlates with increasing dysregulation of glycemic levels. Accordingly one can monitor glycated CD59 levels

over time to determine if glycemic levels of a subject are changing. Changes in relative or absolute glycated CD59 of greater than 0.1% may indicate an abnormality. Preferably, the change in glycated CD59 levels, which indicates an abnormality, is greater than 0.2%, greater than 0.5%, greater than 1.0%, 2.0%, 3.0% , 4.0%, 5.0%, 7.0%, 10%,. 15%, 20%, 25%, 30%, 40%, 50%, or more. Reductions in amounts of glycated CD59 over time may indicated improved glycemic control.

The invention in another aspect provides a diagnostic method to determine the effectiveness of treatments for abnormal glycemic levels. The “evaluation of treatment” as used herein, means the comparison of a subject’s levels of K41-glycated CD59 measured in samples collected from the subject at different sample times, preferably at least one day apart. The preferred time to obtain the second sample from the subject is at least one day after obtaining the first sample, which means the second sample is obtained at any time following the day of the first sample collection, preferably at least 12, 18, 24, 36, 48 or more hours after the time of first sample collection.

The comparison of levels of K41-glycated CD59 in two or more samples, taken on different days, is a measure of level of the subject’s glycemic control and allows evaluation of the treatment to regulate blood sugar levels. The comparison of a subject’s levels of K41-glycated CD59 measured in samples obtained on different days provides a measure of glycemic control to determine the effectiveness of any treatment to regulate blood sugar levels.

As will be appreciated by those of ordinary skill in the art, the evaluation of the treatment also may be based upon an evaluation of the symptoms or clinical end-points of the associated disease, such as the vascular complications of diabetes. Thus, the methods of the invention also provide for determining the regression, progression or onset of a condition which is characterized by abnormal levels of glycated protein, including those characterized by abnormal levels of K41-glycated CD59. In some instances, the subjects to which the methods of the invention are applied are already diagnosed as having a particular condition or disease. In other instances, the measurement will represent the diagnosis of the condition or disease. In some instances, the subjects will already be undergoing drug therapy for regulating blood sugar levels, while in other instances the subjects will be without present drug therapy for regulating blood sugar levels.

According to still another aspect of the invention, a method is provided for treating a subject to reduce the risk of a disorder associated with abnormally-high levels of K41-glycated CD59. The method involves selecting and administering to a subject who is known to have an abnormally-high level of K41-glycated CD59, an agent for
5 treating the disorder. Preferably, the agent is an agent for reducing K41-glycated CD59 levels and is administered in an amount effective to reduce K41-glycated CD59 levels.

In this aspect of the invention, the treatments are based upon selecting subjects who have unwanted, elevated levels of K41-glycated CD59. Such subjects may already be receiving a drug for regulating blood sugar levels, but, according to the invention, are
10 now candidates for an elevated level of the drug based upon the presence of the elevated levels of K41-glycated CD59. It may be appropriate according to the invention to alter a therapeutic regimen for a subject, based upon the measurement of the level of K41-glycated 59. This can be understood in connection with treatment of diabetics.

Diabetics are treated in at least three different ways. Some diabetics are treated only
15 with non-drug therapy, such as exercise and diet. Other diabetics are treated with oral drug therapy, but not with insulin which is injected. Finally, some diabetics are treated with insulin or analogs of insulin by injection. According to the present invention, as a result of determining an elevated level of K41-glycated CD59, an individual undergoing only non-drug therapy may be a candidate for drug therapy as a result of the K41-glycated CD59 test. Likewise, a subject receiving only oral drug therapy, may be a
20 candidate for an insulin-based injectable therapy, due to the K41-glycated CD59 test. Finally, a subject may be free of any present treatment but may be a candidate for blood sugar level regulating treatment as a result of the K41-glycated CD59 test. Thus, subjects may be selected and treated with elevated levels of the same drugs or with
25 different therapies as a result of the assays of the invention.

According to the present invention, some of the subjects are free of symptoms otherwise calling for treatment with a particular therapy. This means that absent the K41-glycated CD59 test, the subject would not according to convention as of the date of the filing of the present application have symptoms calling for treatment with a particular
30 therapy. It is only as a result of the measuring the level of K41-glycated CD59 that the subject becomes a candidate for treatment with the therapy.

Drug therapies for regulating blood sugar levels include oral therapies with hypoglycemic agents an/or oral anti-diabetic agents, injectable therapies, and the like. Non-drug therapies for regulating blood sugar level include, but are not limited to, diatetic and/or exercise control measures.

5 Diet and exercise alterations include, but are not limited to, reducing caloric intake, and/or increasing fiber intake, and/or decreasing fat intake, and/or increasing exercise level.

 Oral drug therapies for regulating blood sugar levels include hypoglycemic agents that may include, but are not limited to:

10 Acarbose; Acetohexamide; Chlorpropamide; Darglitazone Sodium: Glimepiride; Glipizide; Glyburide, Repaglinide; Troglitazone; Tolazamide; Tolbutamide.

 Oral drug therapies for regulating blood sugar levels include antidiabetic agents
15 that may include but are not limited to:
Acarbose, Acetohexamide; Buformin; Butoxamine Hydrochloride ; Camiglibose;
Chlorpropamide; Ciglitazone; Englitazone Sodium; Etoformin Hydrochloride;
Gliamilide; Glibornuride; Glicetanile Gliclazide Sodium; Gliflumide; Glipizide;
Glucagon; Glyburide; Glyhexamide; Glymidine Sodium; Glyoctamide; Glyparamide;
20 Insulin; Insulin, Dalanated; Insulin Human; Insulin Human, Isophane; Insulin Human
Zinc; Insulin Human Zinc, Extended; Insulin, Isophane; Insulin Lispro; Insulin, Neutral;
Insulin Zinc; Insulin Zinc, Extended; Insulin Zinc, Prompt; Linoglriride; Linoglriride
Fumarate; Metformin; Methyl Palmoxirate; Palmoxirate Sodium; Pioglitazone
Hydrochloride; Pirogliride Tartrate; Proinsulin Human; Repaglinide; Seglitide Acetate;
25 Tolazamide; Tolbutamide; Tolpyrramide; Troglitazone; Zopolrestat.

 Injectable therapies for regulating blood sugar levels may include, but are not limited to:

Fast-Acting Insulin:

 Insulin Injection: regular insulin; Prompt Insulin Zinc Suspension; Semilente®
30 insulin. These categories include preparations such as: Humalog® Injection; Humulin®
R; Iletin II; Novolin R, Purified Pork Regular Insulin; Velosulin BR Human Insulin
Intermediate-acting Insulin:

Isophane Insulin Suspension: NPH insulin, isophane insulin; Insulin Zinc Suspension Lente® Insulin. These categories include preparations such as: Humulin® L; Humulin® R; Humulin® N NPH; Iletin® II, Lente®; Iletin® II, NPH; Novolin® L, Novolin® N, Purified Pork Lente® insulin, Purified Pork NPH isophane insulin.

5 Intermediate and Rapid -acting Insulin Combinations:

Human Insulin Isophane Suspension/Human Insulin Injection;. This category includes preparations such as: Humulin® 50/50; Humulin® 70/30; Novolin® 70/30 Long-acting Insulin:

10 Protamine Zinc Insulin Suspension; Extended Insulin Zinc Suspension. These categories include preparations such as: Ultralente® Insulin, Humulin® U.

Reducing the risk of a disorder associated with abnormally high levels of K41-glycated CD59 means using treatments and/or medications to reduce K41-glycated CD59 levels, therein reducing, for example, the subject's risk of vascular complications including but not limited to: diabetic nephropathy, diabetic retinopathy, macro-vascular disease, micro-vascular disease, and diabetic neuropathy.

15 In a subject determined to have an abnormally high level of K41-glycated CD59, an effective amount is that amount effective to reduce K41-glycated CD59 levels in the subject. A response can, for example, also be measured by determining the physiological effects of the hypoglycemic, antidiabetic, or insulin composition, such as the decrease of disease symptoms following administration of the hypoglycemic, antidiabetic, or insulin. Other assays will be known to one of ordinary skill in the art and can be employed for measuring the level of the response. The amount of a treatment may be varied for example by increasing or decreasing the amount of a therapeutic composition, by changing the therapeutic composition administered, by changing the route of administration, by changing the dosage timing and so on. The effective amount will vary with the particular condition being treated, the age and physical condition of the subject being treated, the severity of the condition, the duration of the treatment, the nature of the concurrent therapy (if any), the specific route of administration, and the like factors within the knowledge and expertise of the health practitioner. For example, an effective amount can depend upon the degree to which an individual has abnormally elevated levels of K41-glycated CD59.

An "effective amount" of a drug therapy is that amount of a hypoglycemic, antidiabetic, or insulin or insulin analog that alone, or together with further doses, produces the desired response, e.g. reduction of glycemic level or K41-glycated CD59 levels.

5 In the case of treating a particular disease or condition the desired response is inhibiting the progression of the disease or condition. This may involve only slowing the progression of the disease temporarily, although more preferably, it involves halting the progression of the disease permanently. This can be monitored by routine diagnostic methods known to one of ordinary skill in the art for any particular disease. The desired
10 response to treatment of the disease or condition also can be delaying the onset or even preventing the onset of the disease or condition.

Such amounts will depend, of course, on the particular condition being treated, the severity of the condition, the individual patient parameters including age, physical condition, size and weight, the duration of the treatment, the nature of concurrent therapy
15 (if any), the specific route of administration and like factors within the knowledge and expertise of the health practitioner. These factors are well known to those of ordinary skill in the art and can be addressed with no more than routine experimentation. It is generally preferred that a maximum dose of the hypoglycemic, antidiabetic, or insulin composition (alone or in combination with other therapeutic agents) be used, that is, the
20 highest safe dose according to sound medical judgment. It will be understood by those of ordinary skill in the art, however, that a patient may insist upon a lower dose or tolerable dose for medical reasons, psychological reasons or for virtually any other reasons.

The pharmaceutical compositions used in the foregoing methods preferably are
25 sterile and contain an effective amount of hypoglycemic, antidiabetic, or insulin for producing the desired response in a unit of weight or volume suitable for administration to a patient.

The doses of hypoglycemic, antidiabetic, or insulin administered to a subject can be chosen in accordance with different parameters, in particular in accordance with the
30 mode of administration used and the state of the subject. Other factors include the desired period of treatment. In the event that a response in a subject is insufficient at the

initial doses applied, higher doses (or effectively higher doses by a different, more localized delivery route) may be employed to the extent that patient tolerance permits.

Various modes of administration will be known to one of ordinary skill in the art which effectively deliver the hypoglycemic, antidiabetic, or insulin to a desired tissue, cell or bodily fluid. Preferred methods for administering the hypoglycemic and antidiabetic are oral. The preferred method of administering insulin is by injection. Administration includes: topical, intravenous, oral, intracavity, intrathecal, intrasynovial, buccal, sublingual, intranasal, transdermal, intravitreal, subcutaneous, intramuscular and intradermal administration. The invention is not limited by the particular modes of administration disclosed herein. Standard references in the art (e.g., *Remington's Pharmaceutical Sciences*, 18th edition, 1990) provide modes of administration and formulations for delivery of various pharmaceutical preparations and formulations in pharmaceutical carriers. Other protocols which are useful for the administration of hypoglycemic, antidiabetic, or insulin will be known to one of ordinary skill in the art, in which the dose amount, schedule of administration, sites of administration, mode of administration (e.g., intra-organ) and the like vary from those presented herein.

Administration of hypoglycemic, antidiabetic, or insulin to mammals other than humans, e.g. for testing purposes or veterinary therapeutic purposes, is carried out under substantially the same conditions as described above. It will be understood by one of ordinary skill in the art that this invention is applicable to both human and animal diseases which can be treated by hypoglycemic, antidiabetic or insulin. Thus this invention is intended to be used in husbandry and veterinary medicine as well as in human therapeutics.

When administered, the pharmaceutical preparations of the invention are applied in pharmaceutically-acceptable amounts and in pharmaceutically-acceptable compositions. The term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredients. Such preparations may routinely contain salts, buffering agents, preservatives, compatible carriers, and optionally other therapeutic agents. When used in medicine, the salts should be pharmaceutically acceptable, but non-pharmaceutically acceptable salts may conveniently be used to prepare pharmaceutically-acceptable salts thereof and are not excluded from the scope of the invention. Such pharmacologically

and pharmaceutically-acceptable salts include, but are not limited to, those prepared from the following acids: hydrochloric, hydrobromic, sulfuric, nitric, phosphoric, maleic, acetic, salicylic, citric, formic, malonic, succinic, and the like. Also, pharmaceutically-acceptable salts can be prepared as alkaline metal or alkaline earth salts, such as sodium, potassium or calcium salts. Preferred components of the composition are described above in conjunction with the description of the hypoglycemic, antidiabetic, or insulin compositions of the invention.

A hypoglycemic, antidiabetic, or insulin composition may be combined, if desired, with a pharmaceutically-acceptable carrier. The term "pharmaceutically-acceptable carrier" as used herein means one or more compatible solid or liquid fillers, diluents or encapsulating substances which are suitable for administration into a human. The term "carrier" denotes an organic or inorganic ingredient, natural or synthetic, with which the active ingredient is combined to facilitate the application. The components of the pharmaceutical compositions also are capable of being co-mingled with the hypoglycemic, antidiabetic, or insulin, and with each other, in a manner such that there is no interaction which would substantially impair the desired pharmaceutical efficacy.

The pharmaceutical compositions may contain suitable buffering agents, as described above, including: acetate, phosphate, citrate, glycine, borate, carbonate, bicarbonate, hydroxide (and other bases) and pharmaceutically acceptable salts of the foregoing compounds.

The pharmaceutical compositions also may contain, optionally, suitable preservatives, such as: benzalkonium chloride; chlorobutanol; parabens and thimerosal. The pharmaceutical compositions may conveniently be presented in unit dosage form and may be prepared by any of the methods well-known in the art of pharmacy. All methods include the step of bringing the active agent into association with a carrier which constitutes one or more accessory ingredients. In general, the compositions are prepared by uniformly and intimately bringing the active compound into association with a liquid carrier, a finely divided solid carrier, or both, and then, if necessary, shaping the product.

Compositions suitable for oral administration may be presented as discrete units, such as capsules, tablets, lozenges, each containing a predetermined amount of the active

compound. Other compositions include suspensions in aqueous liquids or non-aqueous liquids such as a syrup, elixir or an emulsion.

Compositions suitable for parenteral administration conveniently comprise hypoglycemic, antidiabetic, or insulin. This preparation may be formulated according to known methods using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation also may be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example, as a solution in 1,3-butane diol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or di-glycerides. In addition, fatty acids such as oleic acid may be used in the preparation of injectables. Carrier formulation suitable for oral, subcutaneous, intravenous, intramuscular, etc. administrations can be found in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA.

The application of the invention to a diabetic subject under treatment with an oral blood sugar regulating agent and otherwise free of symptoms calling for any oral blood sugar regulating agent, as used herein means a subject treated with oral blood sugar regulators whose glycemic-control levels appear normal based on standard diagnostic criteria, including but not limited to measurement of glycated hemoglobin levels.

The application of the invention to a diabetic subject under treatment with insulin (including analogs thereof) and otherwise free of symptoms calling for any insulin, as used herein means a subject treated with insulin whose glycemic-control levels appear to be normal based on standard diagnostic criteria, including but not limited to measurement of glycated hemoglobin levels.

Dosages of blood sugar regulating agents are well-known to those of ordinary skill in the art and documented in the literature.

Referring to Fig. 1, a kit according to the invention is shown. The kit includes a package housing a container which contains an agent for determining the level of K41-glycated CD59 in a sample. The kit also includes a control. The kit also may further comprise instructions, as described above. The instructions typically will be

in written form and will provide guidance for carrying-out the assay embodied by the kit and for making a determination based upon that assay.

EXAMPLES

Example I

5 Introduction

To investigate whether *in vitro* glycation of human CD59 inhibits its homologous restriction activity a functional assay of glycated CD59 was performed. A 10% cell suspension of guinea pig erythrocytes (GPE) was incubated with purified CD59 previously exposed for different time intervals to the glycating sugars glucose (0.5M) or
10 ribose (0.5M), or to non-glycating sorbitol (0.5M) to control for the high osmolarity of the medium. The sensitivity of the GPE to human membrane attack complex (MAC) was tested using purified human C5b6 and C7, C8 and C9 (Halperin et al, 1993a) The number of CD59 molecules incorporated per GPE was determined with 125I-CD59 before and after glycation with ribose.

15 Methods

CD59 purification: CD59 from urine was isolated by anion exchange chromatography using a DEAE protein-Pak HR 8 column (Waters Corp., Milford, MA) as described in Davies et al, 1998. CD59 from butanol extracts of RBC and lysates of CHO cells was purified by immunoaffinity chromatography (see: Davies et al, 1998, Hughes et al, 1992,
20 and van den Berg et al, 1993) using the monoclonal rat anti-human CD59 antibody YTH 53.1 (Serotec Inc. Raleigh, NC), as in Fletcher, et al., 1994. Protein concentration was estimated by the micro-BCA protein assay (Pierce Chemical Co. Rockford, IL). Western blots were performed on proteins separated by SDS-PAGE using the Supersignal detection system (Pierce Chemical Co.).

25 Functional Activity Assays: Hemolysis Protection Assay

Activity of purified CD59 and its mutants before and after glycation was determined by a hemolysis protection assay using guinea pig erythrocytes (GPE) exposed to the terminal complement proteins C5b6, C7, C8 and C9 to form the MAC, as described in Benzaquen, et al, 1994 and Halperin et al., 1993b. In this assay, ~ 10⁻³ unit of human
30 C5b6 is required to lyse 50% of a GPE cell suspension (one unit is the amount of C5b6 required to lyse 50% of a human RBC suspension [Halperin et al, 1993a]). The activity

of purified CD59 was defined as the difference between the percent lysis of the unprotected GPE minus the percent lysis of the CD59-protected GPE.

In vitro glycation: Purified CD59 (10-30 µg/ml) was incubated for different time intervals at 37°C in 0.5 M of either reducing monosaccharides (glucose, ribose, or glucose-6-phosphate) or the non-glycating sorbitol to control for potential osmotic effects.

Incorporation of CD59 into GPE: Purified CD59 was iodinated with 125I (NEN® Life Science Products Inc. Boston, MA) using Iodobeads (Pierce Chemical Co.). Iodination of CD59 did not affect either its homologous restriction activity or its sensitivity to glycation-inactivation. After incubation of GPE with 125I-CD59, the cells were washed extensively to remove unbound CD59, lysed and the radioactivity incorporated was measured in a gamma counter. Using standard methods, the number of CD59 molecules incorporated per GPE was calculated from the specific activity of the 125I-CD59 preparation (150cpm/ng) and the hematocrit of the GPE suspension.

Results

Glycation abrogates the homologous restriction activity of human CD59. The results, illustrated in Figure 2, demonstrate that incubation of CD59 with the glyating sugars glucose and ribose significantly reduce the CD59 activity.

Example 2

Introduction

To determine the location of the site that when glyated inactivates the CD59 molecule, replacement by site-directed mutagenesis of either K41 or H44 was performed. Wild type (WT) and CD59 mutants Gln-41 and Gln-44 were expressed in CHO cells and purified by affinity chromatography. The activity of the mutant CD59 molecules was tested in the GPE hemolytic assay before and after glycation with ribose for different time intervals.

Methods

Expression and purification assays: Using standard methods, wild-type and mutant CD59 cDNAs were subcloned into the mammalian expression vector pSVK3 (Amersham Pharmacia Biotech Inc. Piscataway, NJ) and transfected into CHO cells together with the selection marker pBAGE, which confers resistance to puromycin. Expression and functionality of the WT and mutant CD59 in the puromycin resistant

clones were tested by Western blot analysis, immunohistochemistry in non-permeabilized cells using the YTH 53.1 antibody and a Texas Red conjugated secondary antibody, and a functional dye release assay, which measures protection against human complement, as described in Zhao et al., 1991.

- 5 *Site-directed Mutagenesis:* Plasmid pK562-3 containing the cDNA of human CD59 (Philbrick, et al., 1990) was obtained from ATCC. Site-directed mutagenesis to substitute residue K41 or H44 for glutamine (Gln) was performed using the Altered Sites 11 system (Promega Corp. Madison, WI). The mutagenic primers used were 5'-GTC GTT GAA ATT ACA ATG CTC AAA CTG CCA ACA CTT-3' (SEQ ID NO.: 1) for the
- 10 Gln-41 substitution, and 5'-GTT GAA ATT GCA CTG CTC AAA CTT CCA-3' (SEQ ID NO.: 2) for the Gln-44 mutation. Successful mutagenesis was confirmed by sequencing using standard methods.

For description of *CD59 Purification, In vitro Glycation, and Hemolysis Protection Assay*: See Example 1, Methods.

15 **Results**

Replacement by site-directed mutagenesis of either K41 or H44 abolished the sensitivity of human CD59 to glycation-inactivation confirming that the K41-H44 motif constitute a glycationinactivation site. Results of the GPE lysis assay and the CD59 activity assay are illustrated in Figure 3.

20 Example 3

Introduction

- To determine whether glycation of human endothelial cells and human red blood cells (RBC) results in the inactivation of CD59, an experiment was designed to detect whether glycation would render human endothelial cells and human red blood cells
- 25 (RBC) more sensitive to MACmediated growth factor release or MAC-mediated lysis. Both the endothelial cells and the RBC were glycated and exposed to purified terminal complement components C5b6, C7, C8, and C9 to form the MAC and the mitogenic activity in the conditioned medium was measured.

Methods

- 30 *Glycation of RBC:* A 10% RBC suspension was incubated with or without ribose (50mM) for 48 hours at room temperature, followed by 20 min incubation with cyanoborohydride (20mM). Cell volume was then adjusted by the nystatin procedure

(described in Halperin, et al, 1987) and the osmotic fragility of the cell suspension was measured by standard procedures. To ascertain that differential sensitivity to lysis was not caused by an increased osmotic fragility of RBC exposed to glycating agents, RBC were only used when average cell volume and osmotic fragility were similar to that of control cells.

Glycation of HUVEC: Confluent second passage HUVEC were incubated with or without ribose (50mM) for 24hr (37°C, 5% CO₂) followed by 20min incubation with cyanoborohydride (20mM). Cells were then exposed to purified terminal complement proteins to form the MAC, aliquots of conditioned media separated to test for mitogenic activity in indicator quiescent 3T3 cells, as described in Benzaquen et al, 1994.

RBC and HUVEC Analysis

To avoid confounding effects of colloid osmotic swelling caused by the glycating agent, cell volume was adjusted in all cells, glycated and non-glycated prior to assaying for sensitivity to MAC-induced lysis (Halperin, et al, 1987). After volume adjustment, aliquots of glycated and non-glycated cells were separated for rescue with purified CD59 (4µg), exposure to neutralizing anti-CD59 antibody (YTH53.1) or both, and then exposed to purified C5b6, C7, C8 and C9 to form the MAC.

Results

The data indicate that glycation of the endothelial cells and RBC makes them more sensitive to MAC-mediated growth factor release and MAC-mediated lysis (see Fig. 4). Results are expressed as the ratio of mitogenic activity released into the conditioned media in the presence or absence of MAC.

Example 4

Introduction

To determine whether glycated CD59 is present in human urine, which indicates that glycation occurs in vivo, urine from a nondiabetic subject was concentrated, fractionated, and tested for the presence of CD59 using an anti-CD59 specific antibody. The CD59 positive fractions were then purified further and probed with an antibody that specifically recognizes the reduced ketoamine of glycated lysine residues and binds to the glycated region of CD59, to determine whether glycated CD59 was present in the sample.

Methods

For details of the *CD59 Purification*, see Example 1, Methods section.
Immunoblotting with Anti-hexitol-lysine Antibody was performed using standard procedures as described in Myint, et al, 1995.

Results

5 The CD59 isolated from a non-diabetic subject was immunoblotted with the anti-hexitol-lysine antibody, which recognizes the glycosylated CD59. The results illustrated in Figure 5 demonstrate that glycosylated CD59 is present in human urine.

Example 5

Introduction

10 Following the identification of glycosylated CD59 in human urine, the levels present in urine were measured to determine whether levels of glycosylated CD59 correlate with levels of glycosylated hemoglobin (HbA1c) in diabetic and nondiabetic subjects. Concentrated human urine from either normoglycemic or hyperglycemic (diabetic) individuals was separated by anion exchange chromatography, and a CD59-positive
15 fractions were immunoprecipitated with a monoclonal anti-CD59 antibody. The antibody was used because it binds to an epitope distant from the K41-K44 putative glycosylation motif on CD59 and was, therefore expected to precipitate both glycosylated and non-glycosylated CD59. An aliquot of the immunoprecipitate was separated by SDS-PAGE and blotted with the monoclonal anti-CD59 antibody YTH53.1.

20 Methods

For detail of *CD59 Purification and Quantification* see Example 1, Methods section.

Measurements of HbA1c were performed using standard procedures at the Joslin Diabetic Center (Boston, MA).

25 For details of *Immunoblotting with Anti-hexitol-lysine Antibody*, see Example 4, Methods section.

Results

Comparison of the levels of glycosylated CD59 and glycosylated hemoglobin from diabetic and nondiabetic subjects indicate that the levels of glycosylated CD59 correlate with
30 the levels of glycosylated hemoglobin (see Fig. 5).

Table 1. Sequence Information

AMINO ACID SEQUENCES

SEQ ID NO:1 YNKAWKFEHANFNDC

5

NUCLEIC ACID SEQUENCES

SEQ ID NO:2: GTCGTTGAAATTACAATGCTCAAACCTGCCAACACTT

10

SEQ ID NO:3: GTTGAAATTGCACTGCTCAAACCTTCCA

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183

References

Halperin, J.A. et al., *J. Clin. Invest.* 91, 1974-1978 (1993)

Benzaquen, L.R., et al., *J. Exp. Med.* 179, 985-992 (1994)

Fletcher, C.M., et al., *Structure* 2, 185-199 (1994)

5 Myint, T. et al., *Biochim. Biophys. Acta* 1272, 73-79 (1995)

Davies, A., et al., *J. Exp. Med.* 170, 637-654 (1989)

Hughes, T.R., et al., *Biochem. J.*, 284, 169-176 (1992)

van den Berg., C.W., et al., *Immunology* 78, 349-357 (1993)

Philbrick, W.M., et al., *Eur. J. Immunol.* 20, 87-92 (1990)

10 Zhao, J., et al., *J. Biol. Chem.* 266, 13418-13422 (1991)

Halperin, J.A., et al., *Blood* 81, 200-205 (1993D)

Halperin, J.A. et al., *J. Clin. Invest* 80, 128-137 (1987)

It will be understood that those of ordinary skill in the art that the foregoing represents only certain embodiments of the invention. It will be understood that

15 equivalents of the foregoing are intended to be encompassed within the following claims.

We Claim: